



परीक्षार्थी द्वारा भरा जावे ↓

परीक्षा का विषय: MATHEMATICS: 1 5 0 ENGLISH  
 विषय कोड: 50  
 परीक्षा का माध्यम: ENGLISH

स्टीकर तीर के निशान ↓ से मिलाकर लगायें

माध्यमिक शिक्षा मण्डल, म.प्र., भोपाल  
 पुस्तिका का क्रमांक: C-23 0159348  
 परीक्षार्थी का रोल नम्बर: 2 3 4 4 3 7 3 6 0  
 शब्दों में: TWO THREE FOUR FOUR THREE SEVEN THREE SIX ZERO

नीचे दिये गये उपाहरण अनुसार रोल नम्बर खोजें।

परीक्षार्थी द्वारा भरा जावे ↓

केंद्राध्यक्ष/सहायक केंद्राध्यक्ष एवं परीक्षक द्वारा भरा जावे ↓

प्रश्न पत्र का सेट: D

क :- परीक्षार्थी का कक्ष क्रमांक 20  
 ख :- परीक्षा का दिनांक 21 03 2023  
 परीक्षा का नाम एवं परीक्षा केंद्र क्रमांक की मुद्रा  
 HSSC EXAM C.No.-442199

पर्यवेक्षक का नाम एवं हस्ताक्षर: G. L. Rajouq  
 केन्द्राध्यक्ष/सहायक केन्द्राध्यक्ष के हस्ताक्षर: [Signature]

परीक्षक एवं उपमुख्य परीक्षक द्वारा भरा जावे ↓

प्रमाणित किया जाता है कि होलो क्राफ्ट स्टीकर क्षतिग्रस्त नहीं पाया गया तथा अन्दर के पृष्ठों के अनुरूप मुख्य पृष्ठ पर अंकों की प्रविष्टि एवं अंकों का योग सही है।

निर्धारित मुद्रा: नाम, पदनाम, मोबाईल नम्बर, परीक्षक क्रमांक एवं पदांकित संस्था के नाम की मुद्रा लगाए।  
 उप मुख्य परीक्षक के हस्ताक्षर एवं निर्धारित मुद्रा: परीक्षक के हस्ताक्षर एवं निर्धारित मुद्रा

658  
 2023, 3. 03. 2023  
 परीक्षक G. L. Rajouq (Signature)  
 M. Jari  
 G. L. Rajouq  
 UMS

परीक्षक एवं उपमुख्य परीक्षक द्वारा भरा जावे ↓

केवल परीक्षक द्वारा भरा जावे।  
 प्रश्न क्रमांक के सम्मुख प्राप्तांकों की प्रविष्टि करें।

| प्रश्न क्रमांक | पृष्ठ क्रमांक | प्राप्तांक (अंकों में)   |
|----------------|---------------|--------------------------|
| 1              |               |                          |
| 2              |               |                          |
| 3              |               |                          |
| 4              |               |                          |
| 5              |               |                          |
| 6              |               |                          |
| 7              |               |                          |
| 8              |               |                          |
| 9              |               |                          |
| 10             |               |                          |
| 11             |               |                          |
| 12             |               |                          |
| 13             |               |                          |
| 14             |               |                          |
| 15             |               |                          |
| 16             |               |                          |
| 17             |               |                          |
| 18             |               |                          |
| 19             |               |                          |
| 20             |               |                          |
| 21             |               |                          |
| 22             |               |                          |
| 23             |               |                          |
| 24             |               |                          |
| 25             |               |                          |
| 26             |               |                          |
| 27             |               |                          |
| 28             |               |                          |
| कुल            | में           | कुल प्राप्तांक अंकों में |



प्रश्न क्र.

solution of Question = 6 (not)

Sol

Given that:-

8482210

Direction Ratios of line are  $-18, 12, -4$ i.e.  $a = -18, b = 12, c = -4$  — (1)

Now, According to question

Direction cosines of a line =  $l$  i.e.

$$l, m, n = ?$$

Now,

$$\sqrt{a^2 + b^2 + c^2} = \sqrt{(-18)^2 + (12)^2 + (-4)^2}$$

$$\Rightarrow \sqrt{a^2 + b^2 + c^2} = \sqrt{324 + 144 + 16}$$

$$\Rightarrow \sqrt{a^2 + b^2 + c^2} = \sqrt{484}$$

$$\Rightarrow \sqrt{a^2 + b^2 + c^2} = 22 \quad \text{--- (2)}$$

we know:-

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}, \quad m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \quad n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

$$\Rightarrow l = \frac{-18}{22}, \quad m = \frac{12}{22}, \quad n = \frac{-4}{22}$$

[From eq (1) &amp; (2)]

$$\Rightarrow l = \frac{-9}{11}, \quad m = \frac{6}{11}, \quad n = \frac{-2}{11}$$

Hence, Direction cosines of a line is  $\left( \frac{-9}{11}, \frac{6}{11}, \frac{-2}{11} \right)$

Ans



प्रश्न क्र.

Solution of Question = 7/04

Sol Given that :-

f(x) = x^2 - 4x + 6 - (1)

To find interval in which function is decreasing. On differentiating eq (1) with respect to 'x'

f'(x) = d/dx (x^2 - 4x + 6)

=> f'(x) = d/dx x^2 - d/dx 4x + d/dx 6 [∵ d/dx f1(x) ± f2(x) = d/dx f1(x) ± d/dx f2(x)]

B=> f'(x) = 2x - 4 + 0 [∵ d/dx x^n = nx^(n-1), d(constant) = 0]

S=> f'(x) = 2x - 4

E=> Now, on putting f'(x) = 0

0 = 2x - 4

=> 4 = 2x

=> 2 = x

x = 2 divides real number in two disjoint interval namely (-∞, 2) and (2, ∞)

Test Table

Table with 4 columns: Interval, Test Value, sign of f'(x), Nature of f(x). Rows for (-∞, 2) and (2, ∞).

Hence, In interval (-∞, 2), f'(x) < 0, so, f(x) is decreasing in the interval [-∞, 2] Ans

### Solution of Question = 8

Sol

Given that :-

$$A = \{1, 2, 3\}, B = \{4, 5, 6, 7\}, f = \{(1, 4), (2, 5), (3, 6)\}$$

To show :-

 $f$  is one-oneShowing :-

Here :-

$$f(1) = 4, f(2) = 5, f(3) = 6$$

∴ Image of all distinct element of  $A$  is distinct. So, by definition of one-one

∴ It is one-one.

B

S

Hence Proved :-

 $f$  is one-one Ans

### Solution of Question = 9 (or)

Sol

To Prove :-

$$\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}; x \in [-1, 1]$$

Proof :-

Let -

$$x = \sin \theta \quad \text{--- (1)}$$

$$\Rightarrow x = \cos \left[ \frac{\pi}{2} - \theta \right] \quad \left[ \because \sin x = \cos \left[ \frac{\pi}{2} - x \right] \right]$$

$$\Rightarrow \cos^{-1} x = \frac{\pi}{2} - \theta$$

$$\Rightarrow \cos^{-1} x + \theta = \frac{\pi}{2}$$

$$\Rightarrow \cos^{-1} x + \sin^{-1} x = \frac{\pi}{2} \quad \left[ \because x = \sin \theta \text{ (From eq 1)} \right]$$

$$\Rightarrow \sin^{-1} x = \theta$$

5



Hence Proved :-

$$\sin^2 x + \cos^2 x = \frac{\pi}{2}$$

Ans

Solution of Question = 10

Sol  $A = \begin{bmatrix} -2 & 3 \\ 1 & 2 \end{bmatrix}$ ,  $B = \begin{bmatrix} -1 & 0 \\ 1 & 2 \end{bmatrix}$  (given)

According to question :-

$$(A+B)' = ?$$

Now,

$$A+B = \begin{bmatrix} -2 & 3 \\ 1 & 2 \end{bmatrix} + \begin{bmatrix} -1 & 0 \\ 1 & 2 \end{bmatrix}$$

$$\Rightarrow A+B = \begin{bmatrix} -2+(-1) & 3+0 \\ 1+1 & 2+2 \end{bmatrix}$$

$$\Rightarrow A+B = \begin{bmatrix} -3 & 3 \\ 2 & 4 \end{bmatrix}$$

Now,

$$(A+B)' = \begin{bmatrix} -3 & 3 \\ 2 & 4 \end{bmatrix}'$$

$$\Rightarrow (A+B)' = \begin{bmatrix} -3 & 2 \\ 3 & 4 \end{bmatrix}$$

Hence

$$(A+B)' = \begin{bmatrix} -3 & 2 \\ 3 & 4 \end{bmatrix}$$

Ans

Solution of Question = 11

Sol Let  $y = x^x$

According to question

$$\frac{dy}{dx} = ?$$

Now, on taking log both side in  $y = x^x$

$$\Rightarrow \log y = \log x^x$$

$$\Rightarrow \log y = x \log x \quad [\because \log m^n = n \log m]$$

On differentiating both side with respect to 'x'

$$\Rightarrow \frac{d \log y}{dx} = \frac{d}{dx} x \log x$$

$$\Rightarrow \frac{1}{y} \frac{dy}{dx} = x \frac{d \log x}{dx} + \log x \frac{dx}{dx}$$

$$\left[ \because \frac{d \log x}{dx} = \frac{1}{x}, \frac{d(f_1(x) \cdot f_2(x))}{dx} = f_1(x) \frac{d f_2(x)}{dx} + f_2(x) \frac{d f_1(x)}{dx} \right]$$

B  
S

$$E \Rightarrow \frac{1}{y} \frac{dy}{dx} = x \left( \frac{1}{x} \right) + \log x (1) \quad \left[ \because \frac{d \log x}{dx} = \frac{1}{x}, \frac{dx}{dx} = 1 \right]$$

$$\Rightarrow \frac{dy}{dx} = [1 + \log x] y$$

$$\Rightarrow \frac{dy}{dx} = [1 + \log x] x^x \quad [\because y = x^x]$$

Hence :-

$$\frac{d x^x}{dx} = \underline{\underline{[1 + \log x] x^x}} \quad \underline{\underline{\text{Ans}}}$$

Solution of Question = 12

Sol

Given that :-

$$f(x) = 12x - 3$$

To show :-

$f(x)$  is increasing on  $\mathbb{R}$

Showing :-

7



Let  $x_1, x_2 \in \mathbb{R}$  such that

$$x_1 < x_2$$

$$\Rightarrow 12x_1 < 12x_2 \quad [\text{on multiplying (12) both side}]$$

$$\Rightarrow 12x_1 - 3 < 12x_2 - 3 \quad [\text{on subtracting (3) both side}]$$

$$\Rightarrow f(x_1) < f(x_2) \quad [ \because f(x) = 12x - 3 ]$$

Here,

$$x_1 < x_2 \Rightarrow f(x_1) < f(x_2)$$

So, by definition of increasing function

We can conclude that :-

$f(x) = 12x - 3$  is increasing on  $\mathbb{R}$  Ans

Solution of Question = 13.

Sol Let :-

$$I = \int_{-1}^1 \sin^5(x) \cos^4(x) dx \quad \text{--- (1)}$$

$I = ?$  (According to question)

Here, let :-

$$f(x) = \sin^5(x) \cos^4(x)$$

Now, consider the value of  $f(x)$  at  $(-x)$

$$\Rightarrow f(-x) = \sin^5(-x) \cos^4(-x)$$

$$\Rightarrow f(-x) = (-\sin(x))^5 \cos^4(x) \quad [ \because \sin(-x) = -\sin(x), \cos(-x) = \cos(x) ]$$

$$\Rightarrow f(-x) = -\sin^5(x) \cos^4(x)$$

$$\Rightarrow f(-x) = -f(x)$$

Hence,  $f(x)$  is an odd function

On taking eq (1)

$$I = \int_{-1}^1 \sin^5(x) \cos^4(x) dx$$

$$\Rightarrow I = 0 \quad [ \because \int_{-a}^a f(x) dx = 0 \text{ if } f(x) \text{ is an odd function} ]$$



प्रश्न क्र.

Hence,

$\int \sin^5 x \cos^4 x dx = 0$

Ans

Solution of Question = 14

Sol

Given that :-

$\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$  ,  $\vec{b} = 2\hat{i} + \hat{k}$

Now,

Projection of vector  $\vec{a}$  on the vector  $\vec{b} = ?$

We know :-

Projection of vector  $\vec{a}$  on the vector  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$

$= \frac{(\hat{i} + 2\hat{j} + 3\hat{k}) \cdot (2\hat{i} + \hat{k})}{\sqrt{(2)^2 + (1)^2}}$

$\because \vec{a} \ \& \ \vec{b}$  are given  $|\vec{r}| = \sqrt{a^2 + b^2 + c^2}$

$= \frac{(1)(2) + (2)(0) + (3)(1)}{\sqrt{4+1}}$

$\because \vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$

$= \frac{2 + 0 + 3}{\sqrt{5}}$

$= \frac{5}{\sqrt{5}}$

$= \sqrt{5}$

Hence,

Projection of  $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$  on the vector

$\vec{b} = 2\hat{i} + \hat{k}$  is  $\boxed{\sqrt{5}}$  Ans

B  
S  
E





Solution of Question = 15 (Or)

Sol

$x = ?$  when  $x(\hat{i} + \hat{j} + \hat{k})$  is a unit vector  
Let  $\vec{r} = x(\hat{i} + \hat{j} + \hat{k})$

Now,

$$|\vec{r}| = \sqrt{(x)^2 + (x)^2 + (x)^2} \quad [\because |\vec{a}| = \sqrt{(a_1)^2 + (a_2)^2 + (a_3)^2}]$$

$$\Rightarrow 1 = \sqrt{x^2 + x^2 + x^2} \quad [\because \vec{r} \text{ is a unit vector i.e. } |\vec{r}| = 1]$$

$$\Rightarrow 1 = \sqrt{3x^2}$$

$$\Rightarrow 1 = \pm x\sqrt{3}$$

$$\Rightarrow \frac{\pm 1}{\sqrt{3}} = x$$

Hence,  $\left( \frac{\pm 1}{\sqrt{3}} \right)$  for which  $\vec{r}$  is a unit vector Ans

Solution of Question = 16 (Or)

Sol

Given that :-

$$\frac{dy}{dx} = \frac{1+y^2}{1+x^2} \quad \text{--- (1)}$$

To find general solution of differential equation

Now,

On separating variables in eq (1)

$$\frac{dy}{1+y^2} = \frac{dx}{1+x^2}$$

$$\Rightarrow \frac{dy}{1+y^2} = \frac{dx}{1+x^2}$$

On integrating both side

$$\Rightarrow \int \frac{1}{1+y^2} dy = \int \frac{1}{1+x^2} dx$$



प्रश्न क्र.

$$\Rightarrow \tan^+ y = \tan^+ x + c \quad [\because \int \frac{dx}{1+x^2} = \tan^+ x, \quad c \text{ is a constant}]$$

$$\Rightarrow \tan^+ y - \tan^+ x = c$$

Hence,

General solution of  $\frac{dy}{dx} = \frac{1+y^2}{1+x^2}$  is

$$\tan^+ y - \tan^+ x = c$$

Ans

Solution of Question = 18

Sol  
B  
S  
E

Let

E : Event that the number drawn is more than 3

F : Event that the number drawn is even

Now,

$$S = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$$

$$\Rightarrow n(S) = 10$$

$$E = \{4, 5, 6, 7, 8, 9, 10\}$$

R(x)

$$\Rightarrow n(E) = 7$$

$$P(E) = \frac{n(E)}{n(S)}$$

$$\Rightarrow P(E) = \frac{7}{10} \quad \text{--- (1)}$$

Now

$$F = \{2, 4, 6, 8, 10\}$$

$$\Rightarrow n(F) = 5$$

$$P(F) = \frac{n(F)}{n(S)}$$



$$\Rightarrow P(F) = \frac{5}{10}$$

Now,

$$E \cap F = \{4, 5, 6, 7, 8, 9, 10\} \cap \{2, 4, 6, 8, 10\}$$

$$\Rightarrow E \cap F = \{4, 6, 8, 10\}$$

$$\Rightarrow n(E \cap F) = 4$$

$$P(E \cap F) = \frac{n(E \cap F)}{n(S)}$$

$$\Rightarrow P(E \cap F) = \frac{4}{10} \quad \text{--- (2)}$$

Now, According to question :-

We have to find the Probability of an even number when it is given that number drawn is more than 3 i.e.

$$P\left(\frac{F}{E}\right) = ?$$

Now, We know :-

$$P\left(\frac{F}{E}\right) = \frac{P(E \cap F)}{P(E)}$$

$$\Rightarrow P\left(\frac{F}{E}\right) = \frac{4/10}{7/10} \quad \text{[from eq (1) & (2)]}$$

$$\Rightarrow P\left(\frac{F}{E}\right) = \frac{4}{7}$$

Hence,

Probability of an even number when it is known that drawn card is more than 3 is

$$\frac{4}{7}$$

ANS




प्रश्न क्र.

Solution of Question = 19

Sol

The given curve  $y = x^2$  is parabola which is symmetric about  $y$ -axis and bounded by  $y = 4$



We have :-

B  
S  
E

$y = x^2$   
 $\Rightarrow \sqrt{y} = x$   
Now,

Take a small strip PM

Area of a small strip =  $|x| dy$

Now,

Required area =  $2 \times$  Area of OABD  
 $= 2 \times \int_0^4 |x| dy$

$= 2 \int_0^4 \sqrt{y} dy$  [from eq 1]

$= 2 \cdot 2 \left[ \frac{y^{3/2}}{3/2} \right]_0^4$  [ $\because \int x^n dx = \frac{x^{n+1}}{n+1}$ ]

~~$= \frac{4}{3} [(4)^{3/2} - (0)^{3/2}]$~~

$= \frac{4}{3} [(4)^{3/2} - (0)^{3/2}]$

[ $\because [F(x)]_a^b = F(b) - F(a)$ ]



$$= \frac{4}{3} [4 \cdot 2 - 0]$$

$$= \frac{4 \cdot 8}{3}$$

$$= \frac{32}{3}$$

$$\Rightarrow \text{Required Area} = \frac{32}{3} \text{ sq. unit}$$

Hence, Area of the region bounded by the curve  $y = x^2$  and the line  $y = 4$  is  $\frac{32}{3}$  sq. unit Ans

Solution of Question = 20 (Or)

Sol Given that :-

$$\vec{r} = \hat{i} + 2\hat{j} - 4\hat{k} + \lambda(2\hat{i} + 3\hat{j} + 6\hat{k}) \quad \text{--- (1)}$$

$$\vec{r} = 3\hat{i} + 3\hat{j} - 5\hat{k} + \mu(2\hat{i} + 3\hat{j} + 6\hat{k}) \quad \text{--- (2)}$$

$\therefore$  Both the lines are parallel to  $2\hat{i} + 3\hat{j} + 6\hat{k}$   
So, both the lines are also parallel

According to question :-

Shortest distance between parallel lines (d) = ?

Now, on comparing eq (1) & (2) with standard form :-

$$\vec{r} = \vec{a}_1 + \lambda(\vec{b}) \quad \text{and} \quad \vec{r} = \vec{a}_2 + \mu(\vec{b})$$

$$\vec{a}_1 = \hat{i} + 2\hat{j} - 4\hat{k}, \quad \vec{a}_2 = 3\hat{i} + 3\hat{j} - 5\hat{k}$$

$$\vec{b} = 2\hat{i} + 3\hat{j} + 6\hat{k}$$

Now,

$$\vec{a}_2 - \vec{a}_1 = (3\hat{i} + 3\hat{j} - 5\hat{k}) - (\hat{i} + 2\hat{j} - 4\hat{k})$$

14



पृष्ठ 14 के अंक

कुल अंक



प्रश्न क्र.

$$\Rightarrow \vec{a}_2 - \vec{a}_1 = 3\hat{i} + 3\hat{j} - 5\hat{k} - \hat{i} - 2\hat{j} + 4\hat{k}$$

$$\Rightarrow \vec{a}_2 - \vec{a}_1 = 3\hat{i} - \hat{i} + 3\hat{j} - 2\hat{j} - 5\hat{k} + 4\hat{k}$$

$$\Rightarrow \vec{a}_2 - \vec{a}_1 = 2\hat{i} + \hat{j} - \hat{k}$$

We know :-

Shortest distance between two parallel line (d) =  $\frac{|\vec{b} \times (\vec{a}_2 - \vec{a}_1)|}{|\vec{b}|}$

$$\text{B} \Rightarrow d = \frac{|(2\hat{i} + 3\hat{j} + 6\hat{k}) \times (2\hat{i} + \hat{j} - \hat{k})|}{\sqrt{(2)^2 + (3)^2 + (6)^2}}$$

S  
E

$$\left[ \because \vec{b} = 2\hat{i} + 3\hat{j} + 6\hat{k}, \vec{a}_2 - \vec{a}_1 = 2\hat{i} + \hat{j} - \hat{k}, \right. \\ \left. |\vec{r}| = \sqrt{a^2 + b^2 + c^2} \right]$$

$$\Rightarrow d = \frac{\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 6 \\ 2 & 1 & -1 \end{vmatrix}}{\sqrt{4+9+36}} \quad \left[ \because \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \right]$$

$$\Rightarrow d = \frac{\hat{i}(-3-6) - \hat{j}(-2-12) + \hat{k}(2-6)}{\sqrt{49}}$$

$$\sqrt{\sqrt{49}}$$

$$\Rightarrow d = \frac{|\hat{i}(-9) - \hat{j}(-14) + \hat{k}(-4)|}{7}$$



क्र.

$$\Rightarrow d = \frac{|-9\hat{i} + 14\hat{j} - 4\hat{k}|}{7}$$

$$\Rightarrow d = \frac{\sqrt{9^2 + 14^2 + (-4)^2}}{7} \quad [\because |\vec{r}| = \sqrt{a^2 + b^2 + c^2}]$$

$$\Rightarrow d = \frac{\sqrt{81 + 196 + 16}}{7}$$

$$\Rightarrow d = \frac{\sqrt{293}}{7}$$

Hence,

shortest distance between parallel lines

is  $\frac{\sqrt{293}}{7}$  unitAns

Solution of Question = 21 (Or)

Sol To Prove:-

$$\begin{vmatrix} a-b-c & 2a & 2a \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix} = (a+b+c)^3$$

$$\text{Let } \Delta = \begin{vmatrix} a-b-c & 2a & 2a \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix}$$

On applying  $R_1 \rightarrow R_1 + (R_2 + R_3)$ 

$$\Rightarrow \Delta = \begin{vmatrix} a-b-c+2b+2c & 2a+b-c-a+2c & 2a+2b+c-a-b \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix}$$



प्रश्न क्र.

$$\Rightarrow \Delta = \begin{vmatrix} a+b+c & a+b+c & a+b+c \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix}$$

$$\Rightarrow \Delta = (a+b+c) \begin{vmatrix} 1 & 1 & 1 \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix} \quad \left[ \text{on taking common } (a+b+c) \text{ from } R_1 \right]$$

$$\Rightarrow \Delta = (a+b+c) \begin{vmatrix} 1 & 1 & 1 \\ 2b-2b & b-c-a-2b & 2b \\ 2c-(c-a-b) & 2c-(c-a-b) & c-a-b \end{vmatrix} \quad \left[ \begin{array}{l} \text{on applying} \\ C_1 \rightarrow C_1 - C_3 \text{ \& } \\ C_2 \rightarrow C_2 - C_3 \end{array} \right]$$

**B**  
**S**  
**E**

$$\Rightarrow \Delta = (a+b+c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & -(a+b+c) & 2b \\ 2c-c & 2c-c+a+b & c-a-b \end{vmatrix}$$

$$\Rightarrow \Delta = (a+b+c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & -(a+b+c) & 2b \\ c+a+b & a+b+c & c-a-b \end{vmatrix}$$

On taking common  $(a+b+c)$  from  $C_1$  and  $C_2$

$$\Rightarrow \Delta = (a+b+c)(a+b+c)(a+b+c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & -1 & 2b \\ 1 & 1 & c-a-b \end{vmatrix}$$

On expanding along  $R_1$

$$\Rightarrow \Delta = (a+b+c)^3 \left( 0 \begin{vmatrix} -1 & 2b \\ 1 & c-a-b \end{vmatrix} - 0 \begin{vmatrix} 0 & 2b \\ 1 & c-a-b \end{vmatrix} + 1 \begin{vmatrix} 0 & -1 \\ 1 & 1 \end{vmatrix} \right)$$

$$\Rightarrow \Delta = (a+b+c)^3 (0 - 0 + 1(0(1) - (-1)(1)))$$

$$\Rightarrow \Delta = (a+b+c)^3 (1(1))$$





$\Rightarrow \Delta = (a+b+c)^3$

Hence Proved :-

|         |         |         |               |
|---------|---------|---------|---------------|
| $a-b-c$ | $2a$    | $2a$    | $= (a+b+c)^3$ |
| $2b$    | $b-c-a$ | $2b$    |               |
| $c$     | $2c$    | $c-a-b$ |               |

Ans

Solution of Question = 22 (Del)

Sol Let  $y = \sin x^2$ ,  $w = x^2$   
 On differentiating both side with respect to  $x^2$

According to question :-  
 Differentiation of  $\sin(x^2)$  with respect to  $(x^2)$   
 = ? i.e.

$\frac{dy}{dw} = ?$

On differentiating  $y$  with respect to 'x'

$\frac{dy}{dx} = \frac{d \sin x^2}{dx}$

$\Rightarrow \frac{dy}{dx} = \cos x^2 \frac{d x^2}{dx}$       [ $\because \frac{d \sin x}{dx} = \cos x$ ]

$\Rightarrow \frac{dy}{dx} = \cos x^2 \cdot 2x$       [ $\because \frac{d x^n}{dx} = n x^{n-1}$ ]

$\Rightarrow \frac{dy}{dx} = 2x \cos x^2$       — ①

Now,

On differentiating 'w' with respect to 'x'

प्रश्न क्र.

$$\frac{dw}{dx} = \frac{d x^2}{dx}$$

$$\Rightarrow \frac{dw}{dx} = 2x \quad \left[ \because \frac{d x^n}{dx} = n x^{n-1} \right]$$

Now, dividing eq (1) by eq (2)

$$\frac{dy}{dx} = \frac{2x \cos(x^2)}{2x}$$

$$\Rightarrow \frac{dy}{dw} = \cos(x^2)$$

B  
S  
E

Hence,

Differentiation of  $\sin(x^2)$  with respect to  $x^2$  is  $\boxed{\cos(x^2)}$

Ans

Solution of Question = 23

Sol

Let :-

$$I = \int_0^{x/2} \cos^5 x$$

$$I = \int_0^x \frac{x \sin x}{1 + \cos^2 x} dx \quad \text{--- (1)}$$

Now,

According to question :-

$$I = ?$$

$$I = \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx$$

$$\Rightarrow I = \int_0^{\pi} \frac{(\pi-x) \sin(\pi-x)}{1 + \cos^2(\pi-x)} dx \quad \left[ \because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$\Rightarrow I = \int_0^{\pi} \frac{(\pi-x) \sin x}{1 + \cos^2(-x)} dx \quad \left[ \because \begin{array}{l} \sin(\pi-x) = \sin x, \\ \cos(\pi-x) = -\cos x \end{array} \right]$$

$$\Rightarrow I = \int_0^{\pi} \frac{(\pi-x) \sin x}{1 + \cos^2 x} dx \quad \text{--- (2)}$$

on adding eq (1) & (2)

$$\Rightarrow I + I = \int_0^{\pi} \frac{(\pi-x) \sin x}{1 + \cos^2 x} dx + \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx$$

$$\Rightarrow 2I = \int_0^{\pi} \frac{\pi \sin x - x \sin x + x \sin x}{1 + \cos^2 x} dx \quad \left[ \because \int f_1(x) dx + \int f_2(x) dx = \int [f_1(x) + f_2(x)] dx \right]$$

$$\Rightarrow 2I = \int_0^{\pi} \frac{\pi \sin x}{1 + \cos^2 x} dx \quad \text{--- (3)}$$

Now,

$$\text{Let } \cos x = t$$

$$\Rightarrow -\sin x dx = dt$$

$$\Rightarrow \sin x dx = -dt$$

If  $x = \pi$ , then

$$\cos \pi = t$$

$$\Rightarrow -1 = t \quad \left[ \because \cos \pi = -1 \right]$$

(20)



+



=



योग

पृष्ठ 20 के अंक

पुस्तक संख्या

प्रश्न क्र.

And if  $x=0$ then,  $\cos(0) = t$ 

$$\Rightarrow 1 = t$$

$$[\because \cos 0 = 1]$$

On using  $\sin x dx = -dt$ ,  $t = -1$  and  $t = 1$   
in eq (3)

$$\Rightarrow 2I = \pi \int_{-1}^1 \frac{-dt}{1+t^2}$$

$$\text{B} \Rightarrow 2I = -\pi \int_1^{-1} \frac{dt}{1+t^2}$$

$$\text{S} \Rightarrow 2I = -\pi \left[ \tan^{-1}(t) \right]_1^{-1} \quad \left[ \because \int \frac{dx}{1+x^2} = \tan^{-1}x \right]$$

$$\Rightarrow 2I = -\pi \left[ \tan^{-1}(-1) - \tan^{-1}(1) \right] \quad \left[ \because \int_a^b f(x) dx = F(b) - F(a) \right]$$

$$\Rightarrow 2I = -\pi \left[ -\tan^{-1}(1) - \tan^{-1}(1) \right] \quad \left[ \because \tan^{-1}(-x) = -\tan^{-1}x \right]$$

$$\Rightarrow 2I = -\pi \left[ -\frac{\pi}{4} - \frac{\pi}{4} \right] \quad \left[ \because \tan^{-1}(1) = \frac{\pi}{4} \right]$$

$$\Rightarrow 2I = -\pi \left[ -\frac{\pi}{2} \right]$$

$$\Rightarrow 2I = \frac{\pi^2}{2}$$

$$\Rightarrow 2I = \frac{\pi^2}{2}$$

$$\Rightarrow I = \frac{\pi^2}{4}$$



Hence,

$$\int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx = \frac{\pi^2}{4}$$

Ans

Answer of Question = 1  
Choose the correct Option

(P) (b) 25/A

(ii) (c) Not Defined

(iii) (b)  $\pm 6$

(iv) (a)  $f$  is one-one onto

(v) (b)  $-\frac{\pi}{3}$

(vi) (b)  $\frac{\pi}{3}$

Answer of Question = 2  
Fill in the Blanks

(i)  $\frac{\pi x}{2} + c$

(ii) 0

$$\frac{e^{\sqrt{x}}}{2\sqrt{x}} \cdot 2\sqrt{x}$$



प्रश्न क्र.

(iii) 0

(iv) 1, 0, 0, 0, 1, 0, 0, 0, 1

(v)  ~~$\frac{e^{\sqrt{x}}}{4\sqrt{x}e^{\sqrt{x}}}$  and  $\frac{\sqrt{e^{\sqrt{x}}}}{4\sqrt{x}}$~~

~~(vi)~~

B (v)  $e^x f(x)$

S (vi)  $\frac{e^{\sqrt{x}}}{4\sqrt{x}e^{\sqrt{x}}}$

(vii)  $10 \pi \text{ cm}^2 / \text{cm}$

Answer of Question = 3  
Match the correct pairs

'A'

'B'

(i) Derivative of  $\sin 2x$  with respect to  $x$

(e)  $2 \cos 2x$

(ii)  $\int \tan x dx$

(a)  $-\log|\cos x| + c$

(iii)  $\int \cot x dx$

(b)  $\log|\sin x| + c$



क्र.

(iv)  $\int \sec x dx$  (g)  $\log |\sec x + \tan x| + c$

(v)  $\int \operatorname{cosec} x dx$  (f)  $\log |\operatorname{cosec} x - \cot x| + c$

(vi) 

|          |          |
|----------|----------|
| $\cos x$ | $\sin x$ |
| $\sin x$ | $\cos x$ |

 (e)  $\cos 2x$

Answer of Question = 4

Answer in one word / sentence

B  
S  
E

(i)  $\sqrt{3}$

(ii) 0.4

(iii) 0

(iv) 126

(v) Universal and Empty relation are sometime called Trivial Relation

(vi)  $\frac{\pi}{4}$

(vii) Column matrix is a matrix which has only 1 column i.e.  $A = [a_{ij}]_{m \times 1}$



प्रश्न क्र.

Answer of Question = 5  
True/False

(i) True

(ii) True

(iii) False

B<sup>(iv)</sup> True

S<sup>(v)</sup> True

E<sup>(vi)</sup> false

Question NO. = 17

is. On the

PAGE ~~(31)~~ (29)